



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

INDEX TO VOLUME 42

New names and final members of new combinations are in **bold face type**.

- Abies concolor*, 13; *grandis*, 14, 23; *lasiocarpa*, 13-16
Acer rubrum, 188; *saccharum*, 214
Acorus Calamus, 185
Acrocladium cuspidatum, 189, 204
Aecidium, 57; **Farameae**, 592; *tuberculatum*, 56-58
Aerobryopsis lanosa, 574; *longissima*, 574
Agalinis acuta, 338; *albida*, 391; *decemloba*, 339; *maritima*, 338; *parvifolia*, 340; *paupercula*, 338; *purpurea*, 337, 392
Agromyza Pruni, 235
Agropyron arizonicum, 638; *albicans*, 639; *andinum*, 633-640; *Bakeri*, 638; *caninum*, 638; *dasystachyum*, 637, 639; *Gmelini*, 639; *latiglume*, 633, 640; *molle*, 637, 638; *Palmeri*, 636; *pseudorepens*, 636, 638; *Richardsonii*, 639; *Scribneri*, 633, 634, 640; *Smithii*, 637, 638; *spicatum*, 638, 640; *tenerum*, 638, 639; *violaceum*, 632, 633, 638, 639
Agrostis alba, 192, 635, 636; *asperifolia*, 631, 635, 636; *aequalivalvis*, 636; *canina*, 631, 636; *humilis*, 636, 639; *hyemalis*, 631, 633, 636; *tenuiculmis*, 636; *variabilis*, 631, 636
Aitonina, 261; *australis*, 278; *eximia*, 261; *extensa*, 261; *Fischeriana*, 261; *fissisquama*, 261; *italica*, 278; *japonica*, 261, 304; *lanigera*, 261, 278; *microcephala*, 261; *subplana*, 261; *valida*, 280
Aligrimmia peruviana, 400
Alisma Plantago-aquatica, 183, 194
Allium canadense, 351; *canadense ovoid-eum*, 351; *canadense robustum*, 352
Allium canadense, New varieties of, 351
Alnus incana, 186, 188; *rugosa*, 186-211; *tenuifolia*, 13, 21-24
Alopecurus aristulatus, 631, 636; *occidentalis*, 631, 635, 636
Althaea, 57; *rosea*, 57, 59
Amblystegium, 72; *varium*, 75
Ambrosia artemisiaefolia, 555; *psilostachya*, 555; *trifida*, 555
Ambrosiaceae, 554
Amelanchier, 520, 524; *intermedia*, 527; *oblongifolia*, 188, 205
Ametron, 121
Ammophila arenaria, 551
Ampomele, 121
Amsinckia intermedia, 39
Anagallis arvensis, 29
Anaethaphia attenuata, 379, 381; *bahamensis*, 380, 382; *Buchii*, 380, 385; *calcicola*, 380, 383; *crassifolia*, 380, 382; *Cowellii*, 379, 381; *cuneifolia*, 383; *ilicifolia*, 379, 380; *intertexta*, 380, 384; *lomensis*, 380, 386; *microcephala*, 380, 385; *mantuensis*, 379, 381; *montana*, 379, 381; *Northropiana*, 380, 385; *obtusifolia*, 380, 384; *oligantha*, 380, 382; *Picardae*, 380, 382, 387; *parvifolia*, 380, 383; *paucifloculosa*, 382; *recurva*, 380, 386; *Rosei*, 380, 385; *Shaferi*, 380, 384; *Wilseni*, 380, 383
Anaethaphia D. Don, The genus, 379
Andromeda glaucophylla, 204
Andropogon chrysocomus, 638; *furcata*, 638; *Hallii*, 638
Anogramma refracta, 85; *schizophylla*, 86
Anomobryum densum, 103; *filiforme*, 402
Anomodon, 72; *rostratus*, 74
Anthoceros laevis, 189
Anthoxanthum odoratum, 192
Anthrocephalus, 259, 260; *italicus*, 260, 278; *megacarpus*, 260; *nepalensis*, 259; *polycarpus*, 260
Apargia autumnalis, 549; *autumnalis pratensis*, 550
Apocynaceae, 34
Apocynum album, 36; *androsaeifolium*, 34; *cannabinum*, 35; *hypericifolium*, 36; *medium*, 35; *pubescens*, 35; *sibiricum*, 36
Arctoa, 95
Arethusa bulbosa, 204
Arisaema triphyllum, 188
Arisarum proboscideum, 664-674; *vulgare*, 663-674
Aristida Fendleriana, 640; *longiseta*, 640
Aroids *Arisarum vulgare* and *Arisarum proboscideum*, On the distribution and correlation of the sexes (staminate and pistillate flowers) of the, 663
Aronia arbutifolia, 525, 533; *atropurpurea*, 525; *nigra*, 525
Arthrocormus Schimperii, 572
 ARTHUR, J. C. New Species of Uredineae—IX, 585

- ARTHUR, J. C., & FROMME, F. D. A new North American Endophyllum, 55
 Asclepiadaceae, 36
Asclepias amplexicaulis, 37; *exaltata*, 37; *incarnata*, 185, 204; *pulchra*, 37; *purpurascens*, 36; *syriaca*, 37; *tuberosa*, 36
Aspidium cristatum, 188; *Thelypteris*, 185-188, 204, 207
Aster concolor, 566; *divaricatus*, 564; *dumosus*, 565, 566; *dumosus cordifolius*, 566; *ericoides*, 567; *lateriflorus*, 567; *multiflorus*, 567, 568; *novi-belgii*, 185, 565; *puniceus*, 188; *patens*, 565; *persaliens*, 565; *Pringlei*, 567; *spectabilis*, 566; *subulatus*, 568; *tenuifolius*, 468; *undulatus*, 565; *vimineus*, 567
Astomum chilense, 393
Aulacomnium palustre, 189, 204
Aureolaria pedicularia, 337
Avena americana, 640; *fatua*, 642; *striata*, 640
Aytonia crenulata, 288; *elongata*, 278; *Evansii*, 262, 277, 278; *intermedia*, 301; *jamaicensis*, 262, 277, 292, 296; *mexicana*, 278; *rupestris*, 259, 277
Badiera cubensis, 495, 496; *domingensis*, 494; *diversifolia*, 495; *Fuertesii*, 495; *heterophylla*, 496; *montana*, 495; *Penaea*, 494; *portoricensis*, 494; *propinqua*, 495; *punctata*, 496; *virgata*, 495, 496
Badiera DC., Further studies of, 494
Baeria, 111-115; *aristata*, 115; *Burkei*, 114; *carnosa*, 113, 114; *chrysostoma*, 112, 113, 115; *debilis*, 112; *Fremontii*, 114, 115; *hirsutula*, 113, 115; *microglossa*, 112; *platycarpa*, 113, 114; *tenella*, 114; *uliginosa*, 114, 115
Baeria and *Lasthenia*, Notes on, 111
Barbula, 97; *angustifolia*, 573; *consanguinea*, 573; *fuscineria*, 394; *fuscoviridis*, 394; *lobayetensis*, 573; *pseudoehrenbergii*, 573; *replicata*, 394
Bartonia iodandra, 32, 33; *paniculata*, 32, 33; *virginica*, 32, 33
Bartramia fragilifolia, 402
Batidaea, 122, 127; *amplissima*, 133; *acalyphacea*, 133; *arizonica*, 127, 130; *cataphracta*, 133; *dacotica*, 132, 133; *elegantula*, 132; *heterodoxa*, 132; *itascica*, 27, 130; *laetissima*, 133; *peramoena*, 132, 134; *Sandbergii*, 133; *subarctica*, 132; *subcordata*, 133; *viburnifolia*, 134
Bazzania trilobata, 189, 209
Beckmannia erucaeformis, 635
Benzoin aestivalis, 188
Berberis haematocarpa, 586
Bermuda, Mosses of, 71
Betula fontinalis, 13, 22, 24, 25; *lutea*, 188, 214; *occidentalis*, 24; *papyrifera*, 22, 24, 25; *Piperi*, 24; *subcordata*, 24
BICKNELL, E. P. The ferns and flowering plants of Nantucket—XIV, 27; —XV 331; —XVI, 549
Bidens, 194; *frondosa*, 185
Bignonia lepidota, 377; *microphylla*, 379; *pentaphylla*, 379; *stans*, 372; *Tabebuia*, 372; *triphylla*, 378
 Bignoniaceae, 342
Bihai acuminata, 316, 319; *adflexa*, 316, 319, 323, 325, 326; *angusta*, 316, 319, 327; *aurantiaca*, 316, 319; *barqueta*, 316, 318, 324; *Bihai*, 315-318; *borinquena*, 316, 318, 324; *borinquena coccinea*, 318-324; *brasiliensis*, 316, 319, 324; *brasiliensis pulverulenta*, 319, 326; *Burchellii*, 316, 319; *Collinsiana*, 316, 317; *conferta*, 316, 317; *Champneiana*, 316, 318, 324; *crassa*, 316, 319; *curtispatha*, 317; *dasyantha*, 316, 318; *densa*, 316, 317, 320, 321; *densiflora*, 316, 319; *distans*, 316, 319; *elongata*, 318; *episcopalis*, 316, 317; *geniculata*, 316, 319, 328; *glaucia*, 316, 319; *Harrisiana*, 316, 320, 328, 329; *hirsuta*, 316, 320, 330; *hirsuta cannoidea*, 320, 329; *humilis*, 316, 318, 325; *imbricata*, 315-321; *latispatha*, 316, 319; *librata*, 316, 319; *lingulata*, 316, 319, 323, 326; *longa*, 316, 317; *marginata*, 316, 317; *Mariae*, 316, 317; *metallica*, 316, 318; *pendula*, 316, 317; *platystachys*, 316, 317; *psittacorum*, 315, 316; *pulverulenta*, 326, 327; *punicea*, 316, 317, 321; *purpurea*, 316, 318; *reticulata*, 316, 317; *revoluta*, 316, 317, 322, 326; *rostrata*, 316, 317; *rutila*, 316, 318; *spissa*, 316, 318; *straminea*, 316, 320, 327; *stricta*, 316, 318, 325; *tortuosa*, 316, 318; *velligera*, 316, 318; *villosa*, 316, 317; *Wagneriana*, 316, 317
Bihai, Some new species and varieties of, 315
Blepharoneuron tricholepis, 638
Boehmeria, 39; *cylindrica*, 188
 Boraginaceae, 39
Borrera, 579
Bossekia, 117-119
Botrydium, 194
Bouteloua oligostachya, 638, 640; *prostrata*, 641
Brachythecium novae-angliae, 189
Brasenia Schreberi, 181
 BRITTON, E. G. Mosses of Bermuda, 71; West Indian mosses—II. Mosses of the Danish West Indies and Virgin Islands, 1
 BRITTON, E. G., & HOLLICK, A. A new American fossil moss, 9
 BRITTON, N. L. Studies of West Indian plants—VI, 365; —VII, 487
Bromus breviaristatus, 638; *Flodmanii*,

- 638; lanatipes, 639; hordeaceus, 642; polyanthus, 632, 638; Porteri, 632, 633, 638, 639; Pumpellianus, 633, 638, 639; secundus, 642; Richardsonii, 632, 633, 638
- Bryum* bimum, 189; blandum, 104; capillare, 74; concavum, 402; Cruegeri, 6, 74; decursivum, 5; dichotomum, 74; **microdecurrens**, 5; ovalifolium, 6; ripense, 6; ventricosum, 189
- Bucegia, 265
- Burrielia, 112; leptalea, 112; microglossa, 112
- Buxus* acuminata, 499; bahamensis, 501; brevipes, 499; citrifolia brevipes, 499; cubana, 498; glomerata, 501; gonoclada, 500; Muelleriana, 498; retusa, 500; subcolumnaris, 498; Wrightii, 502
- Calamagrostis* blanda, 636; canadensis, 185, 186, 192, 204; hyperborea, 635, 636; inexpansa, 636; Langsdorfi, 631, 632, 635, 636, 641; laxiflora, 636; micrantha, 636; montanensis, 638; neglecta, 636; purpurascens, 634, 640; Scribneri, 636; Suksdorfii, 641
- Calla palustris*, 204
- Callicostella* Beccariana, 574; papillata, 574; Belangeriana, 7
- Calliergon cordifolium*, 189
- Callirhoë involucrata*, 55-61
- Caltha palustris*, 188
- Calymperes Afzelii*, 1; Breutelii, 3; Dozyanum, 572; hexagonum, 3; Hookeri, 3; lonchophyllum, 3; moluccense, 572; Richardi, 1, 3
- Campanula* aparinoides, 185; rapunculoides, 349
- Campanulaceae, 349
- Campylopus bermudianus*, 71, 72; incrassatus, 393; **peruvianus**, 393
- Caprifoliaceae, 347
- Capsella* Bursa-pastoris, 661; Heegeri, 661, 662
- Cardiobatus*, 122
- Carex* abdita, 621; acuta, 408, 412, 413, 419; acutina, 407, 412, 413; alata, 605; amphiloba, 617; angustata, 407, 412-415; angustata typica, 408; aperta, 410, 411; aquatilis, 406, 407, 414-418; atherodes, 619; brevior, 615; **bulbostylis**, 617; caespitosa, 417; commutata, 408; crinita, 188; Emoryi, 406, 408, 419; **egregia**, 407, 408, 414; **Egglestonii**, 614; erecta, 410; eurycarpa, 414; **exserta**, 620; festiva, 610, 611; **festivella**, 609, 612; fustucacea, 604, 605, 608; festucacea brevior, 604, festucacea tenera, 606; filifolia, 620; filifolia erostata, 620; filiformis, 185, 198, 199; Goodenovii, 417; gravida, 616; grisea, 617; grisea globosa, 617; Haydeni, 406, 407, 410; Kelvingtoniana, 408; Leavenworthii, 618; laeviconica, 519; **Lunelliana**, 615; lurida, 185, 192; macloviana, 610; milligrana, 419; Muhlenbergii, 618; **onusta**, 618; pauciflora, 204; paupercula, 204; Pseudo-Cyperus, 593; rostrata, 204; **rugosperma**, 621; scoparia, 192; **Sheldonii**, 618; stellulata, 192; stipata, 185, 192; straminea, 603, 604, 605, 611; straminea brevior, 603; straminea festucacea, 608; straminea tenera, 606; stricta, 171, 184-192, 405-417; stricta angustata, 406, 408, 419; stricta curtissima, 406, 415; stricta decora, 410; stricta Emoryi, 419; stricta Haydenii, 410; stricta xerocarpa, 407, 408; strictior, 405-417; tenera, 603-606; tenera erecta, 606; tenuispica, 408; torta, 406; trichocarpa, 619; trisperma Billingsii, 204; tristicha, 408; umbellata, 621; virginiana, 408; virginiana elongata, 419; xerocarpa, 408, 410
- Carex*, Miscellaneous new species of, 609
- Carex*, Notes on, — VIII, 405; — IX, 603
- Carex* straminea Willd. and some of its nearest allies, 603
- Carex* stricta Lam. and its allies, 405
- Carpinus caroliniana*, 193
- Carya cordiformis*, 193
- Cassia clarendonensis*, 388; polyadena, 515
- Castalia odorata*, 171, 180, 181
- Castilleja coccinea*, 192
- Catabrosa aquatica*, 630, 635
- Catalpa bignonioides*, 342; *Catalpa*, 342
- Caulerpa clavifera*, 515
- Centaurium spicatum*, 30
- Cephalanthus occidentalis*, 187, 344
- Cephalozia connivens*, 189; fluitans, 204
- Ceratophyllum demersum*, 179, 180
- Ceropteris, 86
- Chaetochloa glauca*, 642
- Chaetomitrium papillifolium*, 574
- Chamaecrista jamaicensis*, 515
- Chamaecyparis*, 519-530; thyoides, 188
- Chamaedaphne*, 209, 212; calyculata, 201-207
- Chamaemorus*, 123; anglica, 123, 124; norwegica, 123
- Chara* sp., 179, 195, 196
- Chiloscyphus rivularis*, 189
- Chiogenes hispidula*, 204, 207
- Chrysanthemum segetum*, 444, 445
- Chrysophyllum stellatum*, 189
- Chrysopsis falcata*, 558
- Chrysosplenium americanum*, 188
- Cichoriaceae, 549
- Cichorium Intybus*, 549
- Cicuta bulbifera*, 183; maculata, 186
- Cinclidotus australis*, 96
- Cinna arundinacea*, 189; latifolia, 640
- Cladium mariscoides*, 183

- Claopodium hawaiense*, 576; *prionophyllum*, 577
Claytonia megarrhiza, 590
Clethra alnifolia, 187, 188; *hyalina*, 261
Climacium americanum, 189
Clintonia borealis, 210
Coccolobis armata, 365, 366; ***benitensis***, 366, 370; ***brevipes***, 366, 371; *calobotrys*, 367; ***colomensis***, 365, 369; *coriacea*, 365, 367; *costata*, 366, 369; ***Cowellii***, 365, 368; *cubensis*, 365, 369; *diversifolia*, 366, 371; *flavescens*, 367; *floridana*, 368; *geniculata*, 366, 370; *laurifolia*, 365, 368; *leoganensis cordata*, 369; *leoganensis parvifolia*, 368; *microphylla*, 365, 366; *nipensis*, 365, 368; *pallida*, 366, 371; *pilonis*, 365, 367; *praecox*, 366, 370; *reflexa*, 370; *retusa*, 365, 368; *rufescens*, 366, 370; *rufescens longifolia*, 370; ***rumicifolia***, 514; ***Shaferi***, 366, 369; *tenuifolia*, 371; *Uvifera*, 365, 367; ***woodfredensis***, 365, 367; *Wrightii*, 366, 371
Coccolobis in Cuba, The genus, 365
Comarobatia, 123, 124
Compositae, 556
Connecticut, The vegetation of. — IV. Plant societies in lowlands, 169
Conocephalum, 272
Convallaria angustifolium, 248; *bifolia canadensis*, 356; *biflora*, 248, 249, 254; *canaliculata*, 248, 249, 255; *commutata*, 248, 256; *hirta*, 248; *multiflora*, 248, 256; *parviflora*, 248; *pubescens*, 248–257
Convolvulaceae, 37
Convolvulus americanus, 38; *arvensis*, 38; *repens*, 38; *sepium*, 37
Coptis trifolia, 188, 210
Cordiobatus, 126
Cornicularia, 579
Cornus Amomum, 187, 188
Corsinia lamellosa, 278
Coumarouna, 625
Crataegus, 13, 21, 24, 230; *mollis*, 521; *punctata*, 521
Crossidium Rosei, 395
Cuba, The genus *Coccolobis* in, 365; The genus *Guettarda* L. in, 507; The genus *Plumiera* L. in, 503; The genus *Scleria* Berg. in, 487; The genus *Tricera* Sw. in, 497
Cucurbitaceae, 349
Cumbata, 121
Cuscuta Gronovii, 38
Cyanea, 77; ***Larrisonii***, 77
Cyanea, A new Hawaiian, 77
Cyclodictyon, 72; *varians*, 74
Cylactis, 120, 121; *lyncimontana*, 120, 125; *montana*, 120, 125
Cymbalaria Cymbalaria, 334
Cyperus calcicola, 387; *dentatus*, 191; *fuliginus*, 387; *strigosus*, 186, 191, 192
Cypripedium acaule, 210
Dactylis glomerata, 642
Dalibarda, 117; *repens*, 117
Danthonia californica, 631, 636; *intermedia*, 631, 636, 640
Datura Stramonium, 334; *Tatula*, 334
Davallia, 651
Decodon verticillatus, 199, 201
Delphinium, 629
Dentaria diphylla, 193
Deschampsia alpica, 631; *atropurpurea*, 631; *caespitosa*, 631, 636; *curtifolia*, 631, 633; *elongata*, 641; *pungens*, 642
Desmatodon subtrophaceus, 398
Dichelostemma pauciflorum
Dicranella flaccidus, 571; *hawaica*, 576; ***Leibergii***, 571; *longirostris*, 2
Dicranoloma angustiflorum, 93; *Blumei*, 572; *grossialare*, 94
Dicranum aucklandicum, 94; *Billardieri*, 95; *flagellare*, 189; *pumilum*, 95; *ramulosum*, 95; *scoparium*, 190, 209
Dictyophlois reticulata, 675–677
Dictyophlois reticulata, gen. et sp. nov., 675
Didymodon calycinus, 95; ***orientalis***, 572; *subtrophaceus*, 398
Didymoglossum, 652
Dioscorea convolvulacea, 592; *grandifolia*, 592
Dipterostemon, 585, 590; *capitatus*, 589; *pauciflorus*, 588, 589
Dipteryx, 623
Distichlis stricta, 637
Distichophyllum microcarpum, 107; *Osterwaldii*, 574
DIXON, H. N. New and rare Australasian mosses, mostly from Mitten's herbarium, 93
DODGE, B. O. The effect of the host on the morphology of certain species of *Gymnosporangium*, 519
Doellingeria humilis, 568; *umbellata*, 568
Dondia intermedia, 592
Drosera longifolia, 202, 204; *rotundifolia*, 202, 204
Dulichium arundinaceum, 171, 183, 204
Dyctisperma, 121
Eatonia intermedia, 636; *pennsylvanica*, 636, 637
Echium vulgare, 41
Ectropothecium cyperoides, 575; *Moritzii*, 575
Effect of the host on the morphology of certain species of *Gymnosporangium*, The, 519
Eleocharis acicularis, 191, 194; *palustris*, 204; *Robbinsii*, 179; *tenuis*, 185; *thermalis*, 642
Elodea canadensis, 179

- Elodium paludosum*, 189
Elymus ambiguus, 641; *canadensis*, 639; *condensatus*, 638; *glaucus*, 632, 637, 641; *innovatus*, 640; *simplex*, 632, 637, 638; *triticoides*, 638
Encalypta emersa, 400
Endophyllum, 55-58; *Sempervivi*, 55, 59; **tuberculatum**, 58, 59, 61
Endophyllum, A new North American, 55
Epilobium densum, 186; sp., 204
Erigeron annuus, 569; *canadense*, 570; *philadelphicus*, 569; *pulchellus*, 569; *pusillus*, 570; *ramosus*, 569
Eriocaulon septangulare, 191
Eriocoma cuspidata, 641
Eriophorum sp., 204; *virginicum*, 186, 201
Erythraea Pickeringii, 30; *spicata*, 30
Erythronium americanum, 193
Eubaeria, 112
Eucladium, 71; *verticillatum*, 73
Eupatorium aromaticum, 557; *hyssopifolium*, 557; *maculatum*, 556; *perfoliatum*, 186, 556; *pubescens*, 557; *purpureum*, 186, 192, 556; *rotundifolium*, 557; *verbenaefolium*, 557
Eupolypodium, 219, 220
Euthamia graminifolia, 564; *minor*, 564; *tenuifolia*, 564
EVANS, A. W. The genus *Plagiochasma* and its North American species, 259
Evernia, 579
Exostema stenophyllum, 517

Fabronia andina, 402
Fagus grandifolia, 214
Faramea occidentalis, 593
FARWELL, O. A. Notes on Michigan Liliaceae, 351; Notes on the Michigan species of *Polygonatum*, 247
Ferns and flowering plants of Nantucket, The, — XIV, 27; — XV, 331; — XVI, 549
Festuca arizonica, 632, 633, 639; *brachyphylla*, 640; *campestris*, 638, 640; *confinis*, 639; *curtifolia*, 634; *elatior*, 637, 641; *fratercula*, 637, 639; *ingrata*, 632, 633, 639; *Jonesii*, 640; *minutiflora*, 634; *octoflora*, 641; *saximontana*, 633, 634, 639, 640; *Thurberi*, 632, 633, 638, 639; *vallicola*, 635, 636
Fissidens, 72; *adiantoides*, 190; *elegans*, 3; *Garberi*, 72; *involutus*, 572; *Kegelianus*, 2; *minutulus*, 73; *Mitteni*, 572; *palmatus*, 2; *taxifolius*, 73; *Zippelianus*, 572
Floribundaria floribunda, 574
FOERSTE, A. F. *Dictyophlois reticulata* gen. et sp. nov., 675
Fossil moss, A new American, 9
Fraxinus americana, 29; *anomala*, 13; *excelsior*, 30; *nigra*, 188; *pennsylvanica*, 193
FROMME, F. D., & ARTHUR, J. C. A new North American *Endophyllum*, 55
Funaria, 71; *calvescens*, 401; *flavicans*, 74; *hygrometrica*, 74, 401; *macrospora*, 401; *suberecta*, 401
Further studies of *Badiera* DC., 494

Galeopsis Ladanum, 42; *Tetrahit*, 42
Galium Claytoni, 189
Gaylussacia baccata, 204, 210, *dumosa*
Bigeloviana, 204
Gentianaceae, 30
Gentiana crinita, 192
Genus *Coccolobis* in Cuba, The, 365
Genus *Guettarda* L. in Cuba, The, 507
Genus *Plagiochasma* and its North American species, The, 259
Genus *Plumiera* in Cuba, The, 503
Genus *Teloschistes* in North America, The, 579
Genus *Tricera* Sw. in Cuba, The, 497
Georgia pellucida, 190, 210
Gerardia decemloba, 339; *paupercula*, 338; *maritima*, 338; *pedicularia*, 337; *purpurea*, 337; *Skinneriana*, 339
Geum rivale, 186
Glaux maritima, 28; *maritima obtusifolia*, 28
Glecoma hederacea, 42
Glycera canadensis, 185; *nervata*, 192; *Torreyana*, 189
Glyphomitrium Cockerelleae, 9
Gochnatia ilicifolia, 380
Galium Aparine, 344; *circaezans*, 347; *Claytoni*, 345, 346; *Claytoni subbiflorum*, 346; *erectum*, 344; *obtusum*, 347; *palustre*, 347; *pilosum*, 344; *tinctorium*, 346; *trifidum*, 346; *triflorum*, 345
Grammitis fluminensis, 223; *limbata*, 222, 223; *linearis*, 225; *marginella*, 221; *quaerenda*, 225
Graphephorum, 635; *muticum*, 631, 636; *Shearii*, 636; *Wolfii*, 631, 636
Gratiola aurea, 191, 336
GRIGGS, R. F. Some new species and varieties of *Bihai*, 315
Grimmia micro-ovata, 400; *saxatilis*, 400
GROSSENBACHER, J. G. Medullary spots and their cause, 227
Guettarda ambigua, 514; *bracteata*, 512; *brevinodis*, 508, 513; **calcicola**, 508, 513; *calypttrata*, 508, 512; **camagueyensis**, 507, 509; **clarensis**, 507, 510; **Coxiana**, 507, 508; *Combsii*, 508, 514; **crassipes**, 507, 511; **cueroensis**, 507, 509; *densiflora*, 507, 511; *Echinodendron*, 507, 508; *elliptica*, 508, 513; *ferruginea*, 507, 511; *havanensis*, 514; *holocarpa*, 509; *inaequipes*, 508, 513;

- lanuginosa, 507, 511; Lindeniana, 508, 513; macrocarpa, 507, 511; monocarpa, 508, 512; multinervis, 513; reticulata, 513; retusa, 508, 514; rigida, 507, 509; rugosa, 514; scabra, 508, 514; sciaphila, 507, 514; undulata, 507, 509; Valenzuelana, 508, 512; **xanthocarpa**, 507, 510; zygomphlebia, 507, 512
- Guettarda L. in Cuba, The genus, 507
- Gymnoconia, 55; interstitialis, 55, 59, 60
- Gymnogramme haematodes, 84; hirta, 80; refracta, 85; schizophylla, 86; Warcewiczii, 80, 84
- Gymnosporangium Blasdaleanum, 520, 521; biseptatum, 519-542; biseptatum follicolum, 520; clavipes, 519, 523; effusum, 529; Ellisii, 523, 528, 530; fraterum, 520-542; globosum, 525; macropus, 519, 521; nidus-avis, 519
- Gymnosporangium, The effect of the host on the morphology of certain species of, 519
- Gymnostomum Breutelii, 3; senocarpum, 3
- Gynoxys, 380
- Gyroweisia, 72; Barbula, 73; boliviana, 394
- Habenaria blephariglottis, 204
- Hagenia, 579
- HALL, H. H. Notes on Baeria and Lasthenia, 111
- Hamamelis virginiana, 190
- HANCE, R. T. Pollen development and degeneration in Zebrina pendula, with special reference to the chromosomes, 63
- Haplocladium microphyllum, 8, 74
- Haplodontium Jamesoni, 401; seriolum, 401
- HARRIS, J. A. On the distribution and correlation of the sexes (staminate and pistillate flowers) in the inflorescence of aroids Arisarum vulgare and Arisarum proboscideum, 663
- Hedeoma pulegioides, 43
- Hedwigidium imberbe, 402
- Heliconia angustata, 327; barqueta, 324; bicolor, 327; brasiliensis, 330; hirsuta, 330; Pearcei, 330; pulverulenta, 326; robusta, 330; stricta, 325
- Hesperopeuce Mertensiana, 17
- Hibiscus Moscheutos, 431-449; oculi-roseus, 429-449
- Hieracium canadensis, 551; aurantiacum, 552; Gronovii, 552; marianum, 552, 553; scabrum, 552, 591; venosum, 552, 553
- HOLLICK, A., & BRITTON, E. G. A new American fossil moss, 9
- Holodontium, 95
- Homaliodendron microdendron, 574
- Hordeum boreale, 636; jubatum, 631, 636, 638; nodosum, 636
- Houstonia coerulea, 343; longifolia, 344
- HOWE, R. H., JR. The genus Teloschistes in North America, 579
- Hypericum mutilum, 191; virginianum, 185, 204
- Hymenophyllum, 651
- Hymenostomum Breutelii, 1, 3; cubense, 3; **ovale**, 576; senocarpum, 3
- Hymenostylium, 71; curvirostre, 73
- Hyophila commutata, 572; **uliginosa**, 4
- Hypnum Antillarum, 2; Breutelii, 2; neckeroides, 105; polystictum, 107; Schreberi, 210
- Hypochaeris radicata, 549
- Hypopterygium Vriesel, 574
- Idaeobatus, 126, 127
- Ilex verticillata, 187, 188, 205
- Ilysanthes dubia, 336; gratioloides, 336
- Impatiens biflora, 185, 188
- Index to American Botanical literature, 49, 87, 161, 241; 309, 423; 481, 543, 595, 643, 679
- Ionactis linariifolius, 568
- Ipomea purpurea, 37
- Iris versicolor, 185, 186
- Isopterygium, 72; minutirameum, 575; micans, 75; tenerum, 71, 75
- Iva frutescens, 554; ovaria, 554; xanthifolia, 554
- Jacandra **Cowellii**, 392
- Juncus acuminatus, 185; canadensis, 185, effusus, 185, 192
- Jungermannia australis, 278; limbata, 278, 281
- Kalmia angustifolia, 204, 210; latifolia, 190; polifolia, 204, 207
- Koeleria gracilis, 638, 640; flexuosa, 43; mutica, 44 Torreyi, 44; verticillata, 44; virginiana, 44
- Krigia virginica, 549
- Labiatae, 41
- Lacinaria scariosa, 558
- Lactuca canadensis, 551; hirsuta, 551; spicata, 551; virosa, 551
- Lamium amplexicaule, 43
- Lappula Lappula, 39
- Larix laricina, 188; Lyallii, 14, 17; occidentalis, 14, 23, 24
- Lasthenia, 111, 116; conjugens, 114, 115; glaberrima, 116; minima, 116
- Lasthenia, Notes on Baeria and, 111
- Ledum groenlandicum, 204
- Leiberg, Mosses of the Philippine and Hawaiian Islands collected by the late John B., 570
- Lemna, 182

- Lentibulariaceae, 340
 Leontodon erythrospermum, 550; Taraxacum, 550
 Leonurus Cardiaca, 43
 Lepidodendron, 675
 Leptamnium virginianum, 342
 Leptilon canadense, 570; pusillum, 570
 Leptobryum pyriforme, 401; Wilsoni, 401
 Leptodontium gracile, 394
 Leskea gracillima, 402
 Leucobryum, 72; glaucum, 72, 189, 209; javense, 572; Polakowskyi, 2; sanctum, 572; sericeum, 572
 Leucoloma Riedlei, 2
 Libocedrus, 520
 Lichen, 579; chrysophthalmus, 579, 581; flavicans, 580
 Liliaceae, Notes on Michigan, 351
 Lillium canadense, 352, 253; **michiganense**, 353; michiganense **umbelliferum**, 353; michiganense **uniflorum**, 353; **peramoenum**, 354
 Lillium, New species of, 352
 Limonium carolinianum, 29
 Limosella tenuifolia, 336
 Linaria canadensis, 335; Cymbalaria, 334; Linaria, 335; vulgaris, 335; vulgaris peloria, 444, 445
 Linnaea americana, 348
 Lippia myrtifolia, 517
 Liriodendron Tulipifera, 193
 Lithospermum arvense, 40
 Lobelia cardinalis, 188; 349; Dortmanna, 191; inflata, 349
 Lobeliaceae, 349
 Lonicera japonica, 349
 Ludvigia palustris, 194
 Lunularia, 270
 Lycium vulgare, 333
 Lycopodium lucidulum, 188
 Lycopodium americanus, 46, 185; communis, 45; membranaceus, 45; rubellus, 45; sessiliflorus, 46; uniflorus, 45; virginicus, 45
 Lymnanthemum lacunosum, 34
 Lyonia ligustrina, 187, 188
 Lysimachia Nummularia, 27; quadrifolia, 27; terrestris, 27, 185, 186, 191, 204; vulgaris, 27
 MACKENZIE, K. K. Notes on Carex—VIII, 405; —IX, 603
 Macromitrium abbreviatum, 100; erosulum, 97–101; grossirete, 100, 101; ligulifolium, 100; oocarpum, 99; orthostichum, 573; papillifolium, 100; per-papillosum, 101; **Petriei**, 101; prorepens, 97–100; Robinsonii, 573; regescens, 101, 102; semipellucidum, 574; submucronifolium, 99
 Maianthemum canadense, 188, 210, 354, 356; canadense interius, 354, 355
 Makinoa crispata, 276
 Manteia, 121
 Marchantia, 265, 272, 273
 Marrubium vulgare, 42
 MAXON, W. R. The North American species of Psilogramme, 79; Polypodium marginellum and its immediate allies, 219
 Mayepea **cubensis**, 390
 Mecosorus marginellus, 221
 Medullary spots and their cause, 227
 Melampyrum lineare, 340
 Melanobatus, 122, 129; bernardinus, 129; glaucifolius, 129; michiganus, 122; neglectus, 122; nigrissimus, 130
 Melica bulbosa, 636; Porteri, 639, 641; spectabilis, 636; subulata, 637, 641
 Melissa officinalis, 43
 Mentha canadensis, 46; crispa, 47; gentilis, 46; piperita, 46; rotundifolia, 46; spicata, 46
 Menyanthes, 34; trifoliata, 34, 204
 Mertensia, 629
 Micrampelis lobata, 349
 Microbryum, 5
 Mielichhoferia campylothecha, 401
 Mikania scandens, 557
 Mimulus ringens, 185
 Mitchella repens, 344
 Mittenothamnium diminutivum, 7
 Mnium cinclidioides, 189; punctatum, 189
 Monopteryx, 623, 625; angustifolia, 623, 625; **Jahnii**, 624–626; Uaucu, 623, 625
 Monopteryx Spruce, On the characters and relationships of the genus, 623
 Mosses from the west coast of South America, 393
 Mosses, mostly from Mitten's herbarium, New and rare Australasian, 93
 Mosses of Bermuda, 71
 Mosses of the Philippine and Hawaiian Islands collected by the late John B. Leiber, 571
 Mosses, West Indian, II. Mosses of the Danish West Indies and Virgin Islands, 1
 Muhlenbergia aristata, 637; comata, 631, 636; filiculmis, 634, 640; filiformis, 636; gracilis, 633, 639; gracillima, 641; pungens, 641; racemosa, 631, 636; Richardsonis, 636, 638; simplex, 637; squarrosa, 636, 638; subalpina, 633, 634, 639, 640; Thurberi, 641; Wolfii, 631, 637; Wrightii, 636
 Munroa squarrosa, 641
 Myosotis arvensis, 40; palustris, 40; scorpioides, 40
 Myrica carolinensis, 205, 210
 Nabalus serpentarius, 553; trifoliolatus, 553

- Najas flexilis*, 179, 191
 Nantucket, The ferns and flowering plants of,—XIV, 27;—XV, 331;—XIV, 549
Nashia cayensis, 517
Neckera disticha, 7; *jamaicense*, 7
Neckeropsis crinita, 574; *Lepiniana*, 574
Neesiella rupestris, 276
Nemopanthus mucronata, 205
Nepeta Cataria, 42
 New American fossil moss, A, 9
 New and rare Australasian mosses, mostly from Mitten's herbarium, 93
 New Hawaiian *Cyanea*, A, 77
 New North American *Endophyllum*, A, 55
 New species of *Lilium*, 352
 New species of *Uredineae*—IX, 585
 New varieties of *Allium canadense*, 351
Nicandra physalodes, 331
 NICHOLS, G. E. The vegetation of Connecticut—IV. Plant societies in lowlands, 169
 North American species of *Psilogramme*, The, 79
 Notes on *Baeria* and *Lasthenia*, 111
 Notes on *Carex*—VIII, 405;—IX, 603
 Notes on Michigan *Liliaceae*, 351
 Notes on *Rosaceae*—IX, 117;—X, 463
 Notes on the Michigan species of *Polygonatum*, 247
 Notes on *Trichomanes*—I. The identity of *Trichomanes pyxidiferum*, 651
Nymphaea advena, 181, 194
Nymphoides lacunosum, 34, 181
Nyssia, 193; *sylvatica*, 193

Octoblepharum albidum, 2, 572
Oenothera alta, 442; *biennis*, 443; *Cockerei*, 443; *cruciata*, 443; *debilis*, 443; *gigas*, 446; *Hookeri*, 443; *laeta*, 442; *Lamarckiana*, 438, 442; *murinella*, 442; *muricata*, 442; *nanella*, 438-445; *Reynoldsii*, 440, 443; *rubrinervis*, 442; *semialta*, 443; *semigigas*, 447; *subrobustus*, 442
 Oleaceae, 29
Onoclea sensibilis, 188
Onosmodium virginianum, 40
 On the characters and relationships of the genus *Monopteryx* Spruce, 623
 On the distribution and correlation of the sexes (staminate and pistillate flowers) in the inflorescence of the aroids *Arisarum vulgare* and *Arisarum proboscideum*, 663
Oreobatus, 119, 120; *deliciosus*, 119; *neomexicanus*, 119; *rubicundus*, 119; *trilobus*, 120
 Origin of dwarf plants as shown in a sport of *Hibiscus oculiroseus*, The, 429
Orobanchaceae, 342
Orobanche uniflora, 342

Orontium aquaticum, 183
Orthomnium stolonaceum, 574
Oryzopsis asperifolia, 640; *exigua*, 639, 641; *micrantha*, 637, 641
Osmunda cinnamomea, 188; *regalis*, 185, 188
 OSTENFELD, C. H. *Ruppia anomala* sp. nov., an aberrant type of the *Potamogetonaceae*, 659
Ostrya virginiana, 14, 25
Ottonia, 261; *italica*, 278; *rupestris*, 278

Pallavicinia Lyellii, 189
Panax trifolium, 193
Panicularia americana, 630, 635; *borealis*, 635; *elata*, 635, 637, 641; *Holmii*, 637; *nervata*, 631-641; *pauciflora*, 630-641
Panicum agrostoides, 191; *barbipulvinatum*, 641, 642; *capillare*, 642; *pacificum*, 636; *thermale*, 642
Paramena, 122, 126; *Menziesii*, 126
Parmelia, 579; *villosa*, 582
 Pasture trees and shubbery, A study of, 451
Pedicularis canadensis, 340
Pelekium velatum, 575
Pellia epiphylla, 189
Peltandra virginica, 183
Petunia axillaris, 334
Phalaris arundinacea, 635, 636
Phascom Floerkeanum, 5; *sessile*, 4
Philonotis australis, 104; *australis surculigera*, 104; *fontana*, 189; *fragilicaulis*, 402; *sphaerocarpa*, 6; *tenella*, 1, 6
Phleum alpinum, 631, 636; *pratense*, 631, 636
Phragmites Phragmites, 635
Phrygilanthus Sonorae, 587
Phyllothallae, 579
Physalis heterophylla, 331; *peruviana*, 331
Physalodes physalodes, 331
Physcia, 579
Physcomitrium Roseae, 400
 Phytogeographical notes on the Rocky Mountain region—IV, Forests of the Subalpine and Montane Zones, 11;—V, Grasslands of the Subalpine and Montane Zones, 629
Picea albertiana, 14, 22, 24; *canadensis*, 22, 25; *Englemannii*, 13-25; *Parryana*, 13-22; *mariana*, 202, 203; *rubra*, 209
Pinus albicaulis, 14-17; *aristata*, 13-19; *edulis*, 13; *flexilis*, 13-24; *monticola*, 14-24; *Murrayana*, 13-24; *ponderosa*, 22, 24; *ponderosa scopulorum*, 19; *scopulorum*, 13-25; *Strobis*, 190
Pirella cymbifolia, 6
Piriqueta cubensis, 389
 PITTIER, H. On the characters and relationships of the genus *Monopteryx* Spruce, 623
Placothallae, 579

- Plagiochasma*, 259; *abyssinicum*, 262; *Aitonia*, 259, 261, 278; *algericum*, 262, 287; *appendiculatum*, 259-305; *articulatum*, 262, 263, 270-276; *Beccarianum*, 262; *brasiliense*, 262; *chlorocarpum*, 260; *Colsmannianum*, 260; *cordatum*, 259; 305; *crenulatum*, 260, 277, 290-305; *cuneatum*, 247; *dschallanum*, 287; *elongatum*, 262, 277, 278, 281, 285, 286, 299, 305; *eximium*, 276, 305; *fissisquamum*, 261; *intermedium*, 260, 263, 276, 277, 301-305; *italicum*, 260, 262, 269, 278, 280, 284; *jamaicense*, 277, 292-305; *japonicum*, 274, 276, 304, 305; **Landii**, 277, 298-304; *lanigerum*, 278, 287, 305; *limbatum*, 260, 278, 285, 305; *mexicanum*, 260, 277, 278, 286, 287; *muricatum*, 281, 288; *nepalense*, 261, 287; *paradoxum*, 260; *pedicellatum*, 260; *peruvianum*, 259, 287; *pterispermum*, 261; *Rousselianum*, 260, 261; *rupestre*, 261-300; *Schimperi*, 262; *subplanum*, 287; *tenuae*, 261, 287, 288; *validum*, 260, 287; *Wrightii*, 260, 277, 294-304
Plagiochasma and its North American species, The genus, 259
Plagiopodopsis Oederi, 10; **Scudderi**, 9, 10
Platycarpha, 113
Platanus occidentalis, 193
Platyklamys, 315
Plantaginaceae, 342
Plantago aristata, 343; *decipiens*, 343; *halophila*, 343; *lanceolata*, 343; *major*, 343; *Rugelii*, 342
Plumbaginaceae, 29
Plumiera clusioides, 503, 504; *confusa*, 505; *emarginata*, 503, 505; *emarginata sericifolia*, 504; *filifolia*, 503, 504; *jamaicensis*, 506; **lanata**, 503, 504; **nipensis**, 503, 505; *obtusa*, 503, 505; *obtusa laevis*, 505; *obtusa parvifolia*, 505; *sericifolia*, 503, 504; **trinitensis**, 503, 506; **venosa**, 503, 506
Plumiera L. in Cuba, The genus, 503
Pneumaria maritima, 39
Poa alpina, 631, 636; *annua*, 642; *aperta*, 633, 639; *Buckleyana*, 632, 638; *callichroa*, 631; *compressa*, 640; *confusa*, 633, 634, 638; *crocata*, 632-640; *Cusickii*, 633, 634, 640; *epilis*, 633, 634, 639; *Fendleriana*, 633, 639, 640; *interior*, 636; *laevigata*, 633, 638, 639; *leptocoma*, 631, 636; *longiligula*, 632-639; *longipedunculata*, 633, 639; *longipila*, 636; *lucida*, 632, 638; *macroclada*, 639; *nervosa*, 631, 636; *nevadensis*, 633, 638, 640; *occidentalis*, 631; *Olneyi*, 631, 636; *paddensis*, 633, 634, 639, 640; *Pattersonii*, 634; *pratensis*, 192; 631, 636; *pratericola*, 636, 638; *pudica*, 631; *reflexa*, 631, 636; *rupicola*, 634; *Sheldonii*, 633, 639; *tricholepis*, 631, 636; *triflora*, 636; *Vaseyana*, 631; *Wheeleri*, 636
Pogonatum polycarpum, 402
Pogonia ophioglossoides, 185, 204
Pohlia, 6; *elongata*, 103; *leptocarpa*, 102; **novae-seelandiae**, 102; *nutans*, 102; *proliger*, 104; *saxensis*, 574
Pollen development and degeneration in *Zebrina pendula*, with special reference to the chromosomes, 63
Polygonatum, 247; *americanum*, 249; *biflorum*, 247-257; *biflorum ovatum*, 352, 355; *biflorum virginicum*, 252, 254; *boreale*, 249-253; *boreale australe*, 252, 253; *canaliculatum*, 249-255; *canaliculatum americanum*, 252, 256; *caniculatum giganteum*, 252, 256; *commutatum*, 249, 251, 256; *cuneatum*, 247, 249, 251, 253; **ellipticum**, 252, 255; *giganteum*, 249, 251, 256; *hirtum*, 251, 252; *latifolium*, 249, 251, 256; *multiflorum*, 249, 251, 254; *parviflorum*, 251; *pubescens*, 249, 250-253; *pubescens cuneatum*, 252, 253; *virginicum*, 249, 251, 254
Polygonatum, Notes on the Michigan species of, 247
Polygonum arifolium, 188
Polypodium ebeninum, 221, 224; *fluminense*, 223; *gramineum*, 219, 220, 205; **Hessii**, 221-224; *leptopodum*, 220, 222; **limbatum**, 220-224; *marginellum*, 219-225; *nigrolimbium*, 220-223
Polypodium marginellum and its immediate allies, 219
Polypogon monspeliensis, 642
Polytrichum commune, 189, 207
Pontederia cordata, 182, 183
Populus acuminata, 25; *angustifolia*, 21-25; *balsamifera*, 21-24; *Besseyana*, 14, 24; *Fremontii*, 13; *hastata*, 14, 24; *Sargentii*, 25; *tremuloides*, 14, 24, 25; *trichocarpa*, 14, 24; *Wislizeni*, 13
Porotrichum latifolium, 106
Portulaca caulerpoides, 515
Potamogeton, 630; *amplifolius*, 178, 179; *epihydrus*, 181; *heterophyllus*, 178, 179; *lucens*, 179; *natans*, 179; *pectinatus*, 179; *pusillus*, 178; *Richardsonii*, 178; *Robbinsii*, 178
Potentilla fruticosa, 193
Preissia, 265, 271
Primulaceae, 27
Proserpinaca palustris, 183, 194
Prunella pennsylvanica, 42; *vulgaris*, 42
Prunus, 230; *avium*, 231; *domestica*, 231; *melanocarpa*, 21
Psedera quinquefolia, 188
Pseudocrossidium, 396; **apiculatum**, 397; **chilense**, 396; **excavatum**, 397
Pseudoleskeopsis decurvata, 575

- Pseudotsuga mucronata*, 13-24
Psilocarya nitens, 387; **portoricensis**, 387
Psilogramme congesta, 79, 81; **chiapensis**, 79, 81; **domingensis**, 86; **flexuosa**, 85; **glaberrima**, 79-83; **glandulosa**, 82; **haematodes**, 80, 84; **hirta**, 82; **hispidula**, 85; **Orbignyana**, 83; **portoricensis**, 80, 85; **refracta**, 80, 84, 85; **schizophylla**, 86; **Warscewiczii**, 79, 80, 81; **villosula**, 83
Psilogramme, The North American species of, 79
Psoralea, 585; **physodes**, 587; **Purshii**, 587
Psychrobatia, 123, 124
Pterobryum angustifolium, 6
Pterocarpus, 625, 626
Pterodon, 625
Pterogoneurum Roseae, 394
Pterygophyllum distichophylloides, 106
Ptilium Crista-castrensis, 209
Ptilomeris, 115
Puccinellia airoides, 637
Puccinia agnita, 585, 590; *Andropogonis*, 538; **Carnegiana**, 587; **Claytoniata**, 591; **Deschampsiae**, 57; **Dondiae**, 585, 592; **Ellisiana**, 521, 538; **Fraseri**, 585, 591; **Hieracium**, 585, 591; **Lojkojana**, 590; **malvacearum**, 60; **nodosa**, 588, 589; **tumamocensis**, 585, 589; **valida**, 585, 591
Puccinosira, 58
Purdiaea microphylla, 389; **Shaferi**, 389; **velutina**, 388
Pycnanthemum incanum, 44
Pyrus americana, 209; **melanocarpa**, 187, 205
Quercus bicolor, 93; **Gambellii**, 13; **macrocarpa**, 13, 25; **palustris**, 193; **undulata**, 13
Racelopus pilifer, 574
Ranunculus aquatilis, 180; **septentrionalis**, 186
Reboulia, 271-274; **hemisphaerica**, 288, 293; **maderensis**, 278, 281, 305
Rhacopilum, 72; **spectabile**, 574; **tomentosum**, 74
Rhamnidium cubensis, 516; **ellipticum**, 516; **Shaferi**, 516
Rheedia Hessii, 390
Rhexia virginica, 192
Rhododendron maximum, 188; **viscosum**, 187, 205
Rhus vernix, 187, 205
Rhynchostegium aristatum, 108, 109; **cylindritheca**, 108, 109; **tenuifolium**, 108, 109
Ribes, 228
Ricciella fluitans, 189
Riccocarpus natans, 189
Rock, J. F. A new Hawaiian *Cyanea*, 77
Roestelia Botryapites, 520, 522, 528, 536; **Ellisii**, 536; **transformans**, 525, 529, 536
Rollandia parvifolia, 78
Rosaceae, Notes on,—IX, 117; —X, 463
Rosa carolina, 187
Rubiaceae, 343
Rubacer, 117-120; **tomentosus**, 119; **velutinus**, 119
Rubus aboriginum, 156; **abundus**, 142; **acalyphaceus**, 133; **acaulis**, 124; **acaulis** × **stellatus**, 464; **acaulis** × **pubescens**, 464; **adenotrichos**, 121, 137, 138, 142; **aegopodioides**, 125; **alleghehiensis**, 149-151; **alleghehiensis** × **argutus**, 467; **alnifolius**, 144; **alpinus**, 142, 143; **ambilis**, 153; **americanus**, 120, 125, 149; **amicalis**, 153; **amicalis** × **nigricans**, 476; **amicalis** × **nigrobaccus**, 468; **amicalis** × **pergratus**, 471; **amplior**, 143, 144; **Andrewsianus**, 150, 151, 465; **arcticus**, 121-124; **arcticus** × **pubescens**, 464; **arcticus** × **saxatilis**, 464; **arenicola** × **hispidus**, 473; **arenicola**, 155; **argutus** × **Baileyanus**, 469; **argutus** × **canadensis**, 471; **argutus** × **carpinifolius**, 470; **argutus** × **cuneifolius**, 465; **argutus** × **Ensenii**, 469; **argutus** × **flagellaris**, 474; **argutus** × **hispidus**, 479; **argutus** × **nigricans**, 476; **argutus** × **nigrobaccus**, 466; **argutus** × **procumbens**, 474; **argutus** × **recurvans**, 469; **arizonicus**, 130, 132; **arundelanus**, 152; **Baileyanus**, 147-158; **Baileyanus** × **canadensis**, 472; **Baileyanus** × **carpinifolius**, 473; **Baileyanus** × **Ensenii**, 475; **Baileyanus** × **frondosus**, 473; **Baileyanus** × **hispidus**, 473; **Baileyanus** × **nigricans**, 477; **Baileyanus** × **nigrobaccus**, 472; **Baileyanus** × **procumbens**, 472; **bernardinus**, 129; **betulifolius**, 151; **Boyntoni**, 155; **Brainerdi**, 147-158; **Brainerdi** × **vermontanus**, 471; **canadensis**, 125-157; **canadensis** × **hispidus**, 479; **canadensis** × **nigricans**, 478; **canadensis** × **nigrobaccus**, 467; **canadensis** × **plicatifolius**, 474; **canadensis** × **Randii**, 471; **canadensis** × **recurvans**, 471; **canadensis** × **sativus**, 466; **carolinianus**, 135; **carpinifolius**, 160; **carpinifolius** × **floridus**, 470; **castoreus**, 464; **Chamaemorus**, 123; **coriifolius**, 139, 140, 141; **costaricanus**, 136, 137; **cubitans**, 479; **cuneifolius**, 147; **cuneifolius** × **floridus**, 466; **cuneifolius** × **lucidus**, 466; **cuneifolius** × **trivalis**, 466; **cyamosus**, 154; **domingensis**, 145; **durus**, 144; **Eastwoodianus**, 146; **Egglestonii**, 132, 135, 136; **elegantulus**, 153; **ellipticus**, 121, 127; **Ensenii**, 157; **Ensenii** × **flagellaris**, 475; **Ensenii** × **frondosus**, 470; **Ensenii** × **hispidus**, 475; **Ensenii** ×

- procumbens, 475; eriocarpus, 128; fagifolius, 121, 139, 143; ferrugineus, 136; flagellaris, 157; flagellaris \times nigricans, 477; flavinanus, 472; floribundus, 142; floricomus, 151; floridus, 151; florulentus, 144, 145; Fockeanus, 124; franciscanus, 126; frondiscentis, 466, 469; frondosus, 147-155; frondosus \times hispidus, 471; frondosus \times nigrobaccus, 467; frondosus \times procumbens, 470; fruticosus, 147, 151; geophilus, 157; glabratus, 126; glandicaulis, 469; glaucifolius, 129; glaucus, 127, 128; Grisebachii, 144; Groutianus, 158; Gunnianus, 124; guyanensis, 141, 143; Helleri, 146; hesperus, 125, 129; heterophyllus, 157; heterophyllus \times nigrobaccus, 468; heterophyllus \times recurvans, 474; hispidus, 156, 187, 188; hispidus major, 478; hispidus \times nigricans, 478; hispidus \times nigrobaccus, 469; hispidus \times plicatifolius, 474; hispidus \times vermontanus, 478; humifusus, 154, 158; humistratus, 154, 158; idaeus, 121-135; idaeus aculeatissimus, 122, 132; idaeus \times phoenicolasius, 464; idaeus \times spectabilis, 464; idaeus \times strigosus, 465; idaeus \times ursinus, 464; invisus, 155; irasuensis, 137, 141; jacens, 478; jamaicensis, 121, 136, 142, 143; Jeckelianus, 152; juncus, 476; laciniatus, 147, 149; laxus, 142; lasiococcus, 123, 124; leucodermis, 122, 129, 130; leucodermis \times melanolasius, 465; Liebmanni, 139; Linkianus, 147; lucidus, 152, 159, 160; macrogongylus, 140; macropetalus, 145, 146, 147; macropetalus \times viburnifolius, 465; melanolasius, 132-135; melanotrachys, 130, 132; Menziesii, 126, 145; Millspaughii, 153; miser, 138; multiformis, 472; multiformis delicatior, 474; multispinus, 474; moluccanus, 118; mucronatus, 125; myriacanthus, 147; neglectus, 465; Nelsonii, 143; neomexicanus, 119; nigerrimus, 129; nigricans, 153, 158, 159; nigricans \times nigrobaccus, 468; nigricans \times plicatifolius, 473; nigricans \times procumbens, 477; nigricans \times Randii, 477; nigricans \times recurvans, 476; nigricans \times sativus, 466; nigricans \times vermontanus, 477; nigrobaccus, 147-156; nigrobaccus \times pergratus, 468; nigrobaccus \times Randii, 468; nigrobaccus \times plicatifolius, 473; nigrobaccus \times vermontanus, 469; nivalis, 122, 126; nutkanus, 119; occidentalis grandiflorus, 122; occidentalis, 125-130; occidentalis \times procumbens, 465; occidentalis \times rosaeifolius, 464; occidentalis \times strigosus, 465; odoratus, 117, 119; oligospermus, 154; ostryifolius, 469; Palmeri, 143; pacificus, 126; parvifolius, 118; parviflorus, 118, 119; Paxii, 464; peculiaris, 476; pedatus, 123, 124; peramoenus, 134, 135; pergratus, 152; persistens, 159, 470; philadelphicus, 152; philyrophyllus, 139; phoenicolasius, 127; pistillatus, 124; Pittieri, 141; plicatifolius, 148-157; poliophyllus, 138, 141; Pringlei, 122, 128; procumbens, 148-158, 465; procumbens \times recurvans, 475; recurvicaulis, 475; recurvicaulis inarmatus, 475; pubescens, 120-130; pumilis, 126; Randii, 150, 151, 154; recurvans, 152, 467; Roezlii, 119; Rossbergianus, 471; rosaeifolius, 122, 126; rhodophyllus, 155; rubrissetus, 160; sativus, 147; saxatilis, 121, 125; sapidus, 139-144; scandens, 121, 138, 139, 143; Schiedeianus, 140; semisetosus, 476; setosus, 158, 159; Smithii, 141; spectabilis, 122, 126, 145, 464; spectabilis \times subarcticus, 464; spectabilis \times viburnifolius, 464; stellatus, 121, 123, 124; strigosus, 125-136, 465; subarcticus, 134, 135; subuniflorus, 157; tardatus, 476; tiliaceus, 139; tiliæifolius, 139; triflorus, 120, 125, 188; trichomallus, 136; trilobus guatemalensis, 120; trifrons, 478, 479; trivialis, 154-160; Tuerckheimii, 144; ursinus, 123, 126, 145, 146; Uhdeanus, 141; urticifolius, 136; Toxicodendron, 188; transmontanus, 121, 125, 130; Verae-Crucis, 137; vermontanus, 153, 158, 159; vermontanus viridifolius, 477; viburnifolius, 134, 135; villosus, 148-157; vitifolius, 146; vulcanicola, 143
- Rupinia australis*, 278; elongata, 278; intermedia, 301; italica, 278; lichenoides, 277; limbata, 278; mexicana, 278; rupestris, 278
- Rupia anomala*, 659; maritima, 659-662; maritima brevirostris, 660
- Rupia anomala* sp. nov., an aberrant type of the Potamogetonaceae, 659
- RYDBERG, P. A. Notes on Rosaceae—I, X, 117; —X, 463; Phytogeographical notes on the Rocky Mountain region —IV, Forests of the subalpine and montane zones, 11; —V, Grasslands of the subalpine and montane zones, 629
- Rynchospora alba*, 201, 204; **borinquensis**, 387; glauca, 387; glomerata, 192
- Sabbatia campanulata*, 30-32; gracilis, 30, 31; stellaris, 31
- Sabina monosperma*, 13; scopulorum, 13, 20, 25; utahensis, 13
- Sagittaria latifolia*, 183
- Salix*, 230; discolor, 187; Lyallii, 24; Mackenziana, 24; Scouleriana, 24; sericea, 187; sessilifolia, 24

- Salomonina biflora*, 253; *commutata*, 254, 255
Salvia officinalis, 43
Sambucus canadensis, 188, 347
Samolus floribundus, 27
Sanguisorba canadensis, 192
Savastana odorata, 637, 641
Sarracenia purpurea, 202
Saxifraga pennsylvanica, 186
Scheuchzeria palustris, 204
Schoenus secans, 489
Schwalbea americana, 340
Scirpus cyperinus, 186; *validus*, 181, 182
Scleria arundinacea, 489; *bracteata*, 491; *bracteata angustata*, 491; *Baldwinii*, 488, 494; *catalinae*, 487, 489; *ciliata*, 487, 490; *Curtissii*, 488, 490; *cubensis*, 488, 489, 492; *debilis*, 489; *dictyocarpa*, 489; *distans*, 488, 492; *Eggersiana*, 491; *elata*, 490; *Elliottii*, 490; *foliosa*, 491; *filiformis*, 493; *gracilis*, 488, 493; *Grisebachii*, 488, 492; *havanensis*, 488, 492; *hemitaphra*, 488; *hirtella*, 488, 492, 493; *interrupta*, 493; *lacustris*, 487, 489; *Liebmanni*, 488, 493; *lithosperma*, 488, 493; *luzulaeformis*, 493; *melaleuca*, 487, 490; *microcarpa*, 488, 491; *microcarpa foliosa*, 491; *mitis*, 488, 491; *nutans*, 493; *Ottonis*, 490; *pauciflora effusa*, 490; *phylloptera*, 488, 491; *pilosissima*, 488, 491; *pinetorum*, 488, 492; *porphyrorrhiza*, 488, 492; *pratensis*, 490; *pterota*, 487, 490; *reflexa*, 489; *reticularis pubescens*, 489; *secans*, 487, 489; *scindens*, 487, 490; *setacea*, 487, 488; *setulosa-ciliata*, 487, 490; *stereorrhiza*, 488, 491; *stereorrhiza major*, 491; *tenella*, 492, 493; *Torreyana*, 488; *verticillata*, 488, 493; *Wrightiana*, 487, 490
Scleria Berg. in Cuba, The genus, 487
 SCOTT, E. L. A study of pasture trees and shrubbery, 451
Scrophularia leporella, 335; *occidentalis*, 336
 Scrophulariaceae, 334
Scutellaria galericulata, 42, 186; *lateriflora*, 42
Sedgwickia hemisphaerica, 278
Selaginella apus, 192
 Selmorition, 121
Sematophyllum, 72; *admistum*, 7; *adnatum*, 75; *hyalinum*, 575; *sericifolium*, 7
 Senecio, 629
Sericocarpus asteroides, 564; *linifolius*, 564
Setiscapella cleistogama, 341; *subulata*, 340, 341
Sherardia arvensis, 347
Sicyos angulatus, 349
Sidalcea, 57; *candida*, 57, 61; *malvaeflora*, 59; *neo-mexicana*, 57, 59
Sigillaria, 675
Sisyrinchium arenicola, 343
Sitanion elymoides, 638, 639; *lanceolatum*, 640; *molle*, 639; *montanum*, 639; *rigidum*, 634, 640
Sium cicutaefolium, 183, 194
 SLOSSON, M. Notes on *Trichomanes*—I. The identity of *Trichomanes pyxidiferum* L., 651
Smilacina canadensis, 355; *canadensis ovalis*, 355, 356; *canadensis trifolia*, 356; *trifolia*, 204
 Solanaceae, 331
Solanum Dulcamara, 333; *nigrum*, 331, 332, 333; *nigrum Dillenii*, 333; *peregrinum*, 332
Solidago, 39; *aestivalis*, 561, 562; *altissima*, 563; *aspera*, 560; *asperula*, 560; *bicolor*, 559; *caesia*, 558; *canadensis*, 563; *Elliottii*, 562; *juncea*, 562, 563; *neglecta*, 186, 562; *nemoralis*, 563; *odora*, 559; *patula*, 188; *puberula*, 559; *rugosa*, 560, 561; *rugosa* × *sempervirens*, 560; *sempervirens*, 560; *uniligulata*, 562
 Some new species and varieties of *Bihai*, 315
Sonchus arvensis, 550; *asper*, 551; *oleraceus*, 550
Sparganium, 630; *americanum androcladum*, 183; *eurycarpum*, 183; *minimum*, 179
Spartina gracilis, 637, 642
Specularia perfoliata, 349
Sphagnum, 72, 189, 196–207; *cuspidatum*, 72; *cymbifolium*, 72; *macrophyllum*, 206; *magellanicum*, 72; *medium*, 72; *trinitense*, 72
Spiraea latifolia, 187; *tomentosa*, 187
Spiridens longifolius, 574
Spirodela, 182
Sporobolus airoides, 638; *asperifolius*, 638, 641; *chryptandrus*, 638; *confusa*, 641; *cryptandrus*, 638, 641
Stachys hyssopifolia, 43
Stenochlamys, 315, 316
Stenolobium stans, 372
Sterophyllum leucostegum, 2, 7
Stigmaria ficoides, 675
Stipa comata, 638; *Elmeri*, 638; *Lettermannii*, 640; *minor*, 632–639; *Nelsonii*, 633, 638, 639; *Porteri*, 637; *Richardsonii*, 638, 640, 641; *Scribneri*, 639; *Tweedyi*, 632, 638; *Vaseyi*, 639, 640; *viridula*, 638–640; *Williamsii*, 638
 STOUT, A. B. The origin of dwarf plants as shown in a sport of *Hibiscus oculi-roseus*, 429
Streblotrichum, 96
 Study of pasture trees and shrubbery, A, 451

- Studies of West Indian plants —VI, 365; —VII, 487
- Symphoricarpos *Symphoricarpos*, 349; *vulgaris*, 349
- Symphytum *aspermum*, 40
- Symplocarpus *foetidus*, 189
- Synthyrisma *humifusum*, 642
- Syringa *vulgaris*, 29
- Syrhropodon, 72; *albovaginatus*, 572; *flavescens*, 3; *floridanus*, 73; *Muelleri*, 572; *subtubulosus*, 572
- Tabebuia *acrophylla*, 378; *actinophylla*, 377; *angustata*, 376; *arenicola*, 375; *arimaensis*, 374; *bahamensis*, 379; *Berterii*, 377; *bibracteolata*, 378; *Brooksiana*, 372, 373; *Buchii*, 377; *calicola*, 373; *crassifolia*, 376; *Curtissii*, 375; *domingensis*, 378; *geronensis*, 375; *heterophylla*, 378; *jamaicensis*, 376; *lepidota*, 377; *moaensis*, 374; *myrtifolia*, 378; *pachyphylla*, 373; *pinetorum*, 374; *platyantha*, 379; *revoluta*, 378; *Shaferi*, 373; *Sauvallei*, 377; *trinitensis*, 373
- Tabebuia Gomez, Additional species of, 372
- Taeniostrobos, 315, 316
- Taraxacum *Dens-leonis*, 550; *erythro-spermum*, 550; *officinale*, 550
- Targionia *hypophylla*, 265, 288
- Taxithelium *alare*, 575; *instratum*, 575; *planum*, 2, 7; *polystictum*, 107
- Taxus *brevifolia*, 14, 24
- Tecoma *acrophylla*, 378; *actinophyllum*, 377; *bahamensis*, 379; *Berterii*, 377; *Brittonii*, 379; *Brittonii decussata*, 379; *Buchii*, 377; *bibracteolata*, 378; *domingensis*, 378; *heptophylla*, 372; *lepidophylla*, 377; *lepidota*, 377; *lepidota myrtifolia*, 378; *Leucoxydon*, 372, 378; *myrtifolia*, 378; *platyantha*, 379; *revoluta*, 378; *sanguinea*, 377; *stans*, 372
- Teloschistes *chrysophthalmus*, 580, 581; *chrysophthalmus flavicans*, 580; *flavicans*, 579, 580; *villosus*, 579–583
- Teloschistes in North America, The genus, 579
- Teucrium *littorale*, 41; *occidentalis*, 41
- Thalesia *uniflora*, 342
- Thalictrum *polygamum*, 186, 189, 192
- Thamniun *arbusculans*, 105, 106; *baculiferum*, 104; *australe*, 106; *latifolium elongatum*, 106
- Thuidium, 72; *delicatulum*, 190; *minutulum*, 74; *plumulosum*, 575
- Thuja *plicata*, 14, 23
- Thymus *Serpyllum*, 44
- Tornabenia, 579
- Tortula, 72; *agraria*, 1, 5, 74; *confusa*, 398; *Kunzeana*, 398; *limensis*, 398; *melanocarpa*, 71, 73; *minuscule*, 399; *muralis*, 399; *verticillata*, 73
- Tragopogon *pratensis*, 550
- Trematodon *capillipes*, 571
- Tricera *acuminata*, 497, 499; *brevipes*, 497, 499; *crassifolia*, 497, 499; *cubana*, 497, 498; *bahamensis*, 497, 501; *fasciculata*, 499; *flaviramea*, 497, 499; *foliosa*, 498, 503; *glomerata*, 497, 501; *gonoclada*, 497, 500; *laevigata*, 498; *Leoni*, 498, 502; *marginalis*, 497, 498; *microphylla*, 498, 502; *Muelleriana*, 497, 498; *retusa*, 497, 500; *revoluta*, 498, 502; *rotundifolia*, 497, 500; *Shaferi*, 497, 500; *subcolumnaris*, 498; *vaccinioides*, 497, 501; *Vahllei*, 502
- Tricera Sw. in Cuba, The genus, 497
- Trichachne *saccharatum*, 633, 639
- Trichocolea *tomentella*, 190
- Trichomanes, 651, 652; *crispum*, 651; *hymenophylloides*, 655, 658; *leptophyllum*, 655; *membranaceum*, 651; *pyxidiferum*, 652–658
- Trichomanes, Notes on,—I. The identity of *Trichomanes pyxidiferum*, 651
- Trichosteleum *hamatum*, 575; *papillatum*, 575
- Trichostema *dichotomum*, 41
- Trichostomum *bermudanum*, 71, 73; *Cockaynii*, 96; *orientale*, 572
- Trientalis *americana*, 28
- Triosteum *perfoliatum*, 348
- Trisetum, 635; *majus*, 631, 633, 636, 639; *montanum*, 632, 641; *subspicatum*, 631, 633, 639
- Trismegistia *lancifolia*, 575
- Tsuga *canadensis*, 190, 193, 214; *heterophylla*, 14, 23; *Mertensiana*, 14
- Typha *latifolia*, 182
- Undescribed West Indian species, 387, 514
- Ulmus *americana*, 14, 25, 188
- Unifolium *bifolium*, 355–358; *bifomuli canadense*, 356; *bifolium kamtschaticum*, 356; *bifolium monophyllum*, 356; *bifolium ovale*, 356; *bifolium trifolium*, 356
- Unifolium *bifolium*, Notes on, 354
- Uredineae, New species of,—IX, 585
- Uredo *fatiscens*, 593
- Uromyces *abbreviatus*, 585, 587; *ornatipes*, 586; *Psoraleae*, 587
- Uropyxis *Woottoniana*, 585
- Usnea, 579
- Utricularia *clandestina*, 340; *cleistogama*, 341; *cornuta*, 204; *geminiscapa*, 340; *intermedia*, 179, 340; *macrorhiza*, 340; *subulata*, 340; *vulgaris americana*, 340
- Vaccinium, 247; *corymbosum*, 187, 188, 205, 210; *macrocarpon*, 204, 207; *Oxycoccus*, 204

- Vagnera stellata*, 357; *stellata mollis*, 357; *trifolia*, 358; *trifolia bifolia*, 358; *trifolia unifolia*, 358
Vagnera, Notes on, 357
Vallisneria spiralis, 179
Vegetation of Connecticut, The,—IV. Plant societies in lowlands, 169
Veratrum viride, 189
Verbascum Blattaria, 334; *phlomoides*, 334; *Thapsus*, 334
Verbena hastata, 41, 186, 192; *urticifolia*, 41
Verbenaceae, 41
Vernonia noveboracensis, 192
Veronica arvensis, 337; *officinalis*, 336; *peregrina*, 337; *serpyllifolia*, 337; *Teucrium*, 337; *Tournefortii*, 337; *virginica*, 193
Vesicularia campylotheca, 575; *filicuspis*, 575; *Meyeriana*, 575
Viburnum dentatum, 187, 188, 347, 348; *cassinoides*, 205; *Lentago*, 188; *molle*, 348; *venosum*, 347, 348
Vinca minor, 34
Viola cucullata, 189; *pallens*, 189
Webera rupestris, 574
Weisia Breutelii, 3; *edentula*, 3; *longiseta*, 73; *Pabstiana*, 3; *seriocarpa*, 3; *viridula*, 73
West Indian mosses—II. Mosses of the Danish West Indies and Virgin Islands, 1
West Indian plants, Studies of, —365; —487
West Indian species, Undescribed, 387, 514
WILLIAMS, R. S., Mosses from the west coast of South America, 393; Mosses of the Philippine and Hawaiian Islands collected by the late John B. Leiberger, 571
Woodwardia virginica, 186, 202, 204
Xanthium echinatum, 556
Zebrina pendula, 63
Zonotrichia, 196
Zygodon circinatus, 400